

# Alternate steroid sulfation pathways targeted by LC-MS/MS 1 analysis of disulfates. Application to prenatal diagnosis of steroid synthesis disorders

Pozo, Oscar J.; Marcos, Josep ; Khymenets, Olha ; Pranata, Andy; Fitzgerald, Christopher C. ; McLeod, Malcolm D. ; Shackleton, Cedric

DOI:

[10.1530/JME-17-0286](https://doi.org/10.1530/JME-17-0286)

License:

Other (please specify with Rights Statement)

*Document Version*

Peer reviewed version

*Citation for published version (Harvard):*

Pozo, OJ, Marcos, J, Khymenets, O, Pranata, A, Fitzgerald, CC, McLeod, MD & Shackleton, C 2018, 'Alternate steroid sulfation pathways targeted by LC-MS/MS 1 analysis of disulfates. Application to prenatal diagnosis of steroid synthesis disorders', *Journal of Molecular Endocrinology*. <https://doi.org/10.1530/JME-17-0286>

[Link to publication on Research at Birmingham portal](#)

## **Publisher Rights Statement:**

Checked for eligibility: 11/07/2018

This is the accepted manuscript for a forthcoming publication in *Journal of Molecular Endocrinology*.

## **General rights**

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

## **Take down policy**

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact [UBIRA@lists.bham.ac.uk](mailto:UBIRA@lists.bham.ac.uk) providing details and we will remove access to the work immediately and investigate.

**Alternate steroid sulfation pathways targeted by LC-MS/MS analysis of disulfates.  
Application to prenatal diagnosis of steroid synthesis disorders**

Oscar J. Pozo<sup>1</sup>, Josep Marcos<sup>2,3</sup>, Olha Khymenets<sup>1</sup>, Andy Pranata<sup>4</sup>, Christopher C.  
Fitzgerald<sup>4</sup>, Malcolm D. McLeod<sup>4</sup> and Cedric Shackleton<sup>5,6</sup>

<sup>1</sup> Integrative Pharmacology and Systems Neuroscience Group, IMIM, Hospital del  
Mar, Doctor Aiguader 88, Barcelona, Spain.

<sup>2</sup> Department of Experimental and Health Sciences, Universitat Pompeu Fabra,  
Doctor Aiguader 88, 08003 Barcelona, Spain.

<sup>3</sup> Cerba Internacional, Pl. Ramon Llull, 7, 08203 Sabadell, Spain.

<sup>4</sup> Research School of Chemistry, Australian National University, Canberra, ACT 2601,  
Australia.

<sup>5</sup> Institute of Metabolism and Systems Research (IMSR), University of Birmingham,  
College of Medical and Dental Sciences, Birmingham, UK.

<sup>6</sup> UCSF Benioff Children's Hospital Oakland Research Institute, Oakland, California.

**Short title**

Steroid disulfates in prenatal diagnosis

**Keywords**

Steroid bis-sulfates, steroid disulfates, steroid sulfation, prenatal diagnosis, LC-  
MS/MS, steroid metabolomics

**Corresponding author: Oscar J. Pozo**

**Word count of the full article 4200**

## ABSTRACT

The steroid disulfates (*aka* bis-sulfates or bis(sulfates)) are a significant but minor fraction of the urinary steroid metabolome that have not been widely studied because major components are not hydrolyzed by the commercial sulfatases commonly used in steroid metabolomics. In early studies, conjugate fractionation followed by hydrolysis using acidified solvent (solvolysis) was used for the indirect detection of this fraction by GC-MS. This paper describes the application of a specific LC-MS/MS method for the direct identification of disulfates in urine, and their use as markers for the prenatal diagnosis of disorders causing reduced estriol production: STSD (Steroid Sulfatase Deficiency), SLOS (Smith-Lemli-Opitz Syndrome) and PORD (P450 Oxido-Reductase Deficiency). Disulfates were detected by monitoring a constant-ion-loss (CIL) from the molecular di-anion. While focused on disulfates, our methodology included an analysis of intact steroid glucuronides and monosulfates because steroidogenic disorder diagnosis usually requires an examination of the complete steroid profile. In the disorders studied, a few individual steroids (as disulfates) were found particularly informative: pregn-5-ene-3 $\beta$ ,20S-diol, pregn-5-ene-3 $\beta$ ,21-diol (STSD, neonatal PORD) and 5 $\alpha$ -pregnane-3 $\beta$ ,20S-diol (pregnancy PORD). Authentic steroid disulfates were synthesized for use in this study as aid to characterization. Tentative identification of 5 $\xi$ -pregn-7-ene-3 $\xi$ ,20S-diol and 5 $\xi$ -pregn-7-ene-3 $\xi$ ,17,20S-triol disulfates was also obtained in samples from SLOS affected pregnancies. Seven ratios between the detected metabolites were applied to distinguish the three selected disorders from control samples. Our results show the potential of the direct detection of steroid conjugates in the diagnosis of pathologies related with steroid biosynthesis.

## INTRODUCTION

From the earliest days of steroid metabolomics, the principal conjugated forms of steroids (sulfates and glucuronides) have been hydrolyzed prior to analysis, and for decades the instrument of choice for steroid separation and measurement has been GC-MS (Shackleton and Marcos 2006). While this technique remains the gold-standard for steroid profiling, LC-MS/MS has been increasingly adopted because of the simplified sample preparation and speed of analysis, mainly provided by absence of a derivatization step. This is in spite of the poor ionization for fully reduced steroids by electrospray (ESI) (Pozo, et al. 2007). While an advance, this methodology still retains the most time-consuming step of sample preparation, the enzymatic or chemical hydrolysis of conjugates (Gomes, et al. 2009). Hydrolysis itself can take several hours and requires a further solid phase extraction (SPE). Necessary chemical derivatization for GC-MS can also take hours.

Intact steroid conjugates have been analyzed by mass spectrometry since the introduction of particle beam ionization (e.g. Fast Atom Bombardment, FAB) in the 1980s (Shackleton and Straub 1982; Shackleton 1983). Their spectra have dominant deprotonated molecules  $[M-H]^-$  in negative ion mode allowing ease of mass determination. Conjugate analysis was simplified with the introduction of electrospray ionization (ESI) and incorporation of HPLC and MS/MS. Glucuronides can be analyzed in both positive and negative ionization modes by monitoring  $[M+NH_4]^+$  and  $[M-H]^-$  respectively (Fabregat, et al. 2013). In the case of monosulfates, collision-induced-dissociation (CID) of the strong  $[M-H]^-$  ions shows a distinctive hydrogen sulfate ( $HSO_4^-$ ) fragment at  $m/z$  97 (Shackleton 1983; Galuska, et al. 2013). Direct detection of steroid conjugates also circumvents the ionization problems of reduced steroids (Pozo, et al. 2007) as phase II metabolites have readily ionized functionality (i.e. a carboxylic acid in glucuronides and an acidic sulfate ester in sulfates).

While mono-conjugates dominate the sulfate fraction of urinary steroids, it has been known since the 1960s that disulfates (diS, also referred to as bis(sulfates) or bis-sulfates to distinguish them from compounds containing the disulfate ( $\text{S}_2\text{O}_7^{2-}$ ) unit) are significant components of the metabolome (Pasqualini and Jayle 1962; Arcos and Lieberman 1967; Shackleton, et al. 1968a; Shackleton, et al. 1968b; Jänne, et al. 1969). Early studies by GC-MS of separated conjugate fractions showed that, in addition to the classic  $3\beta$ -sulfated steroids, hydroxyls at positions  $16\beta$ -,  $17$ -( $\alpha$  and  $\beta$ ) and  $18$ - in androgens and  $20$ - and  $21$ - in pregnanes were prone to sulfation (Jänne, et al. 1969; Jänne and Vihko 1970; Laatikainen, et al. 1972; Meng and Sjövall 1997).

Since these original studies, disulfates have been a largely ignored component of the metabolome that nevertheless had significant potential to expand the understanding of steroid biosynthetic and metabolic pathways. Given this, we sought to develop LC-MS/MS methodology to target this group. It was found that constant-ion-loss (CIL) of hydrogen sulfate ( $\text{HSO}_4^-$ ) fragment at  $m/z$  97 from the molecular di-anion  $[\text{M}-2\text{H}]^{2-}$  was the most useful reaction to monitor (McLeod, et al. 2017).

The ease of steroid disulfate analysis led us to investigate their use in diagnosis of steroid biosynthetic disorders. One particular area of interest to the authors has been the pre-natal diagnosis of single-gene disorders of estriol (E3) synthesis by urine analysis, of which we have studied three conditions by GC-MS, viz., Steroid Sulfatase Deficiency (STSD), Smith-Lemli-Opitz Syndrome (SLOS, 7-dehydrosterol reductase deficiency) and cytochrome P450 Oxido-Reductase Deficiency (PORD) (Marcos et al. 2009; Shackleton, et al. 2004a; Shackleton et al. 2004b; Arlt et al. 2004; Reisch, et al. 2013; Shackleton et al. 2007). This communication offers our preliminary observations of the disulfated steroids excreted in these disorders at around mid-pregnancy. While focusing on disulfates, selected monosulfates and

103 glucuronides were also included; evaluating the complete steroid profile is crucial to  
104 diagnosing aberrant steroid biosynthesis (Shackleton and Marcos 2006).

105

## MATERIALS AND METHODS

### *Reagents and chemicals*

Steroid starting materials were obtained from Steraloids (Newport, RI, USA). Chemicals and solvents including sulfur trioxide pyridine complex (SO<sub>3</sub>.py), *N,N*-dimethylformamide (DMF) and ammonium formate (HPLC grade) were purchased from Sigma–Aldrich (St Louis, MO, USA). Aqueous ammonia solution (25%), and acetonitrile and formic acid (LC-MS grade) were from Merck (Darmstadt, Germany). MilliQ water was obtained using a Milli-Q purification system (Millipore Ibérica, Barcelona, Spain).

### *Synthesis of reference steroid disulfates*

The qualitative synthesis of steroid disulfates as the ammonium salts was performed as previously described (McLeod, et al. 2017) with small modifications. Briefly, 1 mg of each steroid standard was directly dissolved in a freshly prepared solution of SO<sub>3</sub>.py complex (20 mg, 124 µmol, ~38 eq/steroid or ~19eq/hydroxyl group) in DMF (100 µL) and incubated at room temperature for 72 hours. The success of synthesis was confirmed by analysis of reaction using both LC-MS in scan mode and LC-MS/MS for collision induced dissociation studies. The purification of synthesised disulfates was performed using SPE as previously described (McLeod et al, 2017).

Steroid disulfate reference materials isolated as the corresponding ammonium salts and used in this study included: 5 $\alpha$ -pregnane-3 $\beta$ ,20S-diol bis(sulfate), (3 $\beta$ 5 $\alpha$ PD-diS); 3 $\beta$ ,21-dihydroxypregn-5-en-20-one bis(sulfate), (21-hydroxypregnenolone bis(sulfate), 21OHPreg-diS); androst-5-ene-3 $\beta$ ,17 $\alpha$ -diol bis(sulfate), (5AD(17 $\alpha$ )-diS); androst-5-ene-3 $\beta$ ,17 $\beta$ -diol bis(sulfate), (5AD(17 $\beta$ )-diS); 3 $\beta$ ,16 $\alpha$ -dihydroxyandrost-5-en-17-one bis(sulfate), (16 $\alpha$ -hydroxydehydroepiandrosterone bis(sulfate),

16 $\alpha$ OHDHEA-diS); 3 $\beta$ ,16 $\beta$ -dihydroxyandrost-5-en-17-one bis(sulfate) (16 $\beta$ -hydroxydehydroepiandrosterone bis(sulfate), 16 $\beta$ OHDHEA-diS); pregn-5-ene-3 $\beta$ ,17 $\alpha$ ,20S-triol 3,20 bis(sulfate), (5PT-diS); pregn-5-ene-3 $\alpha$ ,20S-diol bis(sulfate), (5PD-diS); 5 $\beta$ -pregnane-3 $\beta$ ,20S-diol bis(sulfate),; 5 $\beta$ -pregnane-3 $\alpha$ ,20S-diol bis(sulfate),; 5 $\alpha$ -pregnane-3 $\alpha$ ,20S-diol bis(sulfate),; 5 $\beta$ -pregnane-3 $\beta$ ,20R-diol bis(sulfate), 5 $\alpha$ -pregnane-3 $\beta$ ,20R-diol bis(sulfate),; 5 $\beta$ -pregnane-3 $\alpha$ ,20R-diol bis(sulfate),; In this manuscript the IUPAC terms for the 20-hydroxypregnane diastereomers are used, S and R, in some publications often trivialized to  $\alpha$  and  $\beta$ , respectively.

Two reference materials (3 $\beta$ 5 $\alpha$ PD-diS and 21OHPreg-diS), were prepared on larger scale and subjected to characterisation by spectroscopic methods. Experimental details and characterization data for these new compounds, together with copies of the  $^1\text{H}$  NMR,  $^{13}\text{C}$  NMR, and ESI LRMS spectra are available from the authors (MM).

### ***Urine Samples***

One of our laboratories (Children's Hospital Oakland, Dr. Cedric Shackleton) has been the recipient for urine samples from patients with suspected abnormal steroidogenesis in an attempt to characterize the defects. The studies were approved by the Children's Hospital Institutional Review Board (IBR#2010-038)). Many of the samples used in this study were remnants of those sent to the laboratory for investigation of low pregnancy estriol (generally defined as individuals with serum unconjugated estriol < 0.3 MoM, multiples of median). Other samples were from women who had had a previously affected SLOS child or other symptomatic reasons for concern regarding steroidogenesis. The samples have generally been collected between week 16 and 30 of gestation. They have been stored frozen at -20 °C. Eleven STSD samples were analyzed, and six samples from SLOS affected pregnancies. The neonatal PORD samples were collected at 7, 18 and



23 days. Urine samples from unaffected pregnancies were from a collection held by IMIM (Institut Hospital del Mar d'Investigacions Mèdiques, Barcelona). Normal neonatal urine specimens were from a control urine collection at the Institute of Metabolism and Systems Research (IMSR), University of Birmingham UK.

### ***Sample treatment***

Urine extraction was by C18 SPE. Generally, a 2 mL aliquot of urine was passed through a pre-conditioned cartridge. After a washing step with 3 mL water, steroid conjugate analytes were eluted using 2 mL of methanol. After evaporation of a 200  $\mu$ L aliquot of the elution solvent, the extract was reconstituted in 100  $\mu$ L of water and 5  $\mu$ L was injected into the UHPLC-MS/MS system. Stably labelled 17-S $\{^{18}\text{O}\}_3$ -5 $\alpha$ -androstane-3 $\alpha$ ,17 $\beta$ -diol 3,17-bis(sulfate) and 17-S $\{^{18}\text{O}\}_3$ -5 $\alpha$ -androstane-3 $\beta$ ,17 $\beta$ -diol bis (sulfate) were used as internal standards. The labeled sulfate residue was introduced to the steroidal diol mono-sulfate using labelled S $\{^{18}\text{O}\}_3$ .py generated in situ from labelled sulfuric acid (95% atom) and acetic anhydride in pyridine. Experimental details and characterization data for these internal standards, together with copies of the  $^1\text{H}$  NMR,  $^{13}\text{C}$  NMR, and ESI LRMS spectra are available from the authors (MM)

### ***UHPLC-MS/MS analysis***

#### ***Disulfates***

The study was carried out using a triple quadrupole (XEVO TQ-S micro) mass spectrometer equipped with an ESI source and interfaced to an Acquity UPLC system

for the chromatographic separation (all from Waters Associates, Milford, MA, USA). Drying gas as well as nebulizing gas was nitrogen. The desolvation gas flow was set to approximately 1200 L/h, and the cone gas flow was 50 L/h. A cone voltage of 30 V and a capillary voltage of 0.4 kV were used in negative ionization mode. The nitrogen desolvation temperature was set to 600 °C, and the source temperature was 150 °C.

The UHPLC separation was performed using an Acquity UPLC CSH Phenyl-Hexyl column (2.1 × 100 mm i.d., 1.7 µm) (Waters Associates), at a flow rate of 300 µL/min. Water and acetonitrile:water (9:1) both with formic acid (0.01% v/v) and ammonium formate (25 mM) were selected as mobile phase solvents. A gradient program was used; the percentage of organic solvent was linearly changed as follows: 0 min, 15%; 0.5 min, 15%; 25 min, 30%; 26 min, 100%; 27 min, 100%; 28 min, 15%; 30 min, 15%. The total analysis time was 30 min.

For the constant ion loss (CIL) scan, dwell times of 6 ms and collision energies of 15 eV were selected for each ion transition. Due to the molecular masses of steroid hormones and metabolites (250-400 Da), the precursor ions of disulfates ( $[M-2H]^{2-}$ ) were restricted to the range from  $m/z$  199 to  $m/z$  274. A Selected Reaction Monitoring (SRM) approach containing 75 preselected transitions was used for the simultaneous detection of steroid disulfates. Among them, the transition 228→359 corresponded to the internal standards used in the analysis.

### *Monoconjugates*

While the focus has been on steroid disulfates we have acquired data on steroid monosulfates and glucuronides previously reported as relevant for the studied disorders. Based on previous studies (Gomez, et al. 2014) the product ions at  $m/z$  97 and  $m/z$  75 for sulfates and glucuronides respectively were chosen (Table 1). Exceptions were estriol conjugates due to the influence of the aromatic ring. The

neutral loss of the conjugate (80 Da and 176 Da for sulfates and glucuronides respectively were detected).

### *Quantification*

For this study, accurate quantitative measurements have not been conducted for two reasons: 1) lack of some authentic compounds prevented the determination of relative responses of analyte transitions to internal standard transitions; 2) the urine samples were random “spot” collections and not accurate 24 h collections. Instead, we have determined “diagnostic-ratios” from raw mass spectrometric transition responses. These ratios are of an analyte known to be *overproduced* to one known to be *underproduced* in a particular disorder. Such ratios have long been used in GC-MS analysis (Shackleton and Marcos,2006).

## RESULTS AND DISCUSSION

### *Method development*

This communication applies recent LC-MS/MS studies on steroid disulfate analysis using the constant-ion-loss (CIL) from the di-anionic precursor  $[M-2H]^{2-}$  (McLeod, et al. 2017). The method was developed for untargeted detection, and designed for the analysis of a maximum number of natural disulfates. The use of this precursor ion and the fact that the product ion has a higher  $m/z$  value is unusual for small molecules. Determination of disulfates under these conditions gives clean chromatograms and the main interferences observed in the chromatograms are due to the relatively high natural abundance of the  $^{34}\text{S}$  isotope (4.25%). The transition coming from the  $m/z$  97 loss from an unsaturated ( $\Delta^4$ ,  $\Delta^5$ , etc.)  $\{^{34}\text{S}\}_1$ -disulfate isotope is completely indistinguishable from the one coming from an A-ring reduced steroid disulfate.

To maximize isobaric steroid metabolite separation (e.g. pregnenediol-diS, the pregnenediol-disulfates and the androstenediol-disulfates) in this study, a phenyl-hexyl column with a relatively high amount of ammonium formate (25 mM) was required to obtain sharp and well resolved chromatographic peaks. Column temperature was critical for this purpose with 30 °C determined as optimum. Under these conditions, a 25 min gradient from 15% to 30% of organic solvent provided desired separation (Figure 1A).

Under optimized conditions the elution order of disulfates was dihydroxyandrostanes < dihydroxypregnanones < androstenediols < pregnenediols. In a specific group,  $17\beta$  hydroxysteroid disulfates eluted earlier than their  $17\alpha$ -counterparts and  $20S$ -hydroxysteroid disulfates eluted earlier than their  $20R$  counterparts. Regarding A ring derivatives,  $\Delta^5$  steroid disulfates eluted before

the fully reduced metabolites, the elution order of the reduced steroids being  $3\beta,5\beta < 3\beta,5\alpha < 3\alpha,5\beta < 3\alpha,5\alpha$ . The chromatographic conditions were also able to separate the two estriol glucuronide isomers i.e. the 16-glucuronide and 3-glucuronide. Unfortunately, sulfate and glucuronide conjugates of two useful steroids in PORD diagnosis, androsterone and etiocholanolone, could not be separated under the selected conditions even after increasing the gradient to 1 h (Figure 1B).

### ***Application to prenatal detection of disorders affecting estriol synthesis***

We report preliminary studies to determine whether steroid disulfates in urine can be useful markers in the prenatal detection of disorders affecting estriol synthesis; until now only monoconjugates had been used. The background to this study being that unconjugated serum E3 is frequently measured at mid-pregnancy as a marker for Down's syndrome as part of a test called triple- or quad- marker screening (Haddow, et al. 1994). If results are low the question remains as to the reason, and our original research was directed to diagnosis of Smith-Lemli-Opitz Syndrome (SLOS), the clinically most severe cause of low E3 (Shackleton, et al. 2007). These studies led to investigation of other causes such as STSD and PORD.

Diagnostic ratios are frequently employed in steroid metabolomics and E3 frequently used as denominator. Dominant E3 conjugates are 3- and 16-glucuronides (30% and 60%, respectively) with about 2.5% as monosulfate and estriol-3-glucuronide-16-sulfate (6.5%) (Tikkanen, et al. 1973). We assessed E3 excretion from the measurement of glucuronide and monosulfate conjugates (Table 1).

### ***Steroid sulfatase deficiency (STSD) (OMIM , 308100, location, Xp22.31)***

This X-linked disorder prevents the release of steroid from steroid sulfates. A summary of the biosynthetic pathway leading to estriol is shown in Figure 2, illustrating that inactivity of the enzyme in placenta prevents  $16\alpha$ OHDHEA-S conversion to E3. This fetal  $16\alpha$ OHDHEA-S, androst-5-ene- $3\beta$ , $16\alpha$ , $17\beta$ -triol sulfate (5AT-S) and other steroid sulfates pass through the placenta and mother to be excreted in urine largely unchanged (Taylor and Shackleton, 1979).

STSD urine samples (N=11) and 11 controls were analyzed using the CIL scan method for disulfates complemented with the acquisition of  $16\alpha$ OHDHEA-S and E3 monoconjugates (Table 1). Among the disulfates measured by the CIL method, we found that the response ratio between six of them, namely  $16\alpha$ OHDHEA-diS, 5AD( $17\alpha$ )-diS, 5AD( $17\beta$ )-diS, 5PT-diS, 21OHPreg-diS and 5PD-diS against E3 glucuronide (measured as sum of 3- and 16-glucuronides) was markedly increased in STSD.

Representative chromatograms of a normal pregnancy urine and one with an STSD affected fetus are shown in Figure 3. The ratio values for our normal and STSD data sets are shown in Figure 4A and show all analytes clearly distinguish STSD from normal. Additionally, we used the ratios to evaluate the relative efficacy of each analyte in diagnosis. The best steroid discriminatory ratio would show greatest difference between the lowest steroid sulfate/E3-G ratio value in STSD, and the highest ratio found in controls (Figure 4B). Interestingly, the ratios that gave the greatest differential were  $\Delta^5$  pregnenes; 5PD-diS, 5PT-diS and 21OHPreg-diS, not the  $C_{19}$  steroid sulfates on the direct biosynthetic pathway to E3. Combining ratio data can give an even greater separation for normal and affected; note the combined data for 21OHPreg-diS and 5PD-diS in Figure 4C. Such pregnene metabolites should be incorporated in MS based methodologies for detection of the disorder.

*P450 oxido-reductase deficiency (PORD) (POR OMIM 124015 location: 7q11.23)*

311

312 Several pregnant women carrying PORD fetuses have been studied by GC-MS  
313 (Shackleton, et al. 2004; Reisch, et al. 2013), but for only two were samples available  
314 for this study. Shackleton and co-workers (2004a) deduced that the dominant “feto-  
315 placental” maternal urinary steroid in PORD pregnancies was  $3\beta 5\alpha$ PD-diS. While this  
316 steroid is also present in the disulfate fraction of normal pregnancy urine it is in  
317 much greater amount in PORD affected pregnancies. From its dominance together  
318 with reduced E3, it was concluded that this metabolite is a maternal excretory  
319 product of fetal pregnenolone. An intermediate precursor would be fetal steroid  
320 5PD-diS. Excess pregnenolone and its sulfate are the result of an apparent “block” in  
321 17-hydroxylase/C17-20 lyase secondary to attenuated POR activity (Figure 5). This  
322 block, together with suppressed  $16\alpha$ -hydroxylase (also due to PORD) causes  
323 reduction of fetal  $16\alpha$ OHDHEA-S production leading to low maternal E3 production  
324 and excretion. The precise sequence of reactions from fetal pregnenolone to  
325  $3\beta 5\alpha$ PD-diS, and localities of the conversions (fetal adrenal, liver, placenta and  
326 mother) is yet to be determined. The process is multi-step, probably including  
327 placental  $3\beta$ -desulfation and likely  $3\beta$ -hydroxysteroid dehydrogenase/isomerase. It  
328 has long been known that both  $3\beta 5\alpha$ PD-diS and 5PD-diS are prominent disulfates in  
329 umbilical cord blood (Laatikainen, et al. 1972) so are freely synthesized and  
330 transported in the feto-placental unit. The synthetic sequence for pregnenolone  
331 conversion to urinary metabolites in normal and PORD affected pregnancies and  
332 neonate are shown in Figure 5. Evidence suggests the corresponding conversion of  
333 pregnenolone sulfate to DHEA-S is not an available pathway (Neunzig, et al. 2014;  
334 Sanchez-Guijo, et al. 2016; Rege, et al. 2017).

335 Besides the increased excretion of  $3\beta 5\alpha$ PD-diS, we also observed an increase in the  
336 transitions corresponding to 5PD-diS and  $3\beta, 21$ -dihydroxy- $5\alpha$ -pregnan-20-one  
337 disulfate ( $21\text{OHPreg}3\beta 5\alpha$ -diS), the latter in spite of a likely POR requirement by fetal  
338 21-hydroxylase. However, it should be noted that this fetal enzyme differs from that

339 coded by CYP21A2 required in cortisol synthesis (Guerami et al., 1988, Corsan,  
340 Macdonald and Casey., 1997).

341 In Figure 6 we illustrate the chromatographic profiles of the  $3\beta 5\alpha$ PD-diS, 5PD-diS  
342 and 21OHPreg $3\beta 5\alpha$ -diS and the E3 conjugates in a control (Figure 6A) and affected  
343 pregnancy (Figure 6B). The dominance of the  $3\beta 5\alpha$ PD-diS in the affected  
344 pregnancies is striking. In GC-MS diagnosis of PORD prenatally the ratio of  
345  $3\beta 5\alpha$ PD/E3 was used, i.e. the ratio of principal PORD fetal metabolite to E3, the  
346 conventional feto/maternal metabolite. In Figure 6C are shown ratios for intact  
347 conjugates in PORD and controls. For the denominator (E3) we summed the total of  
348 both glucuronylated forms.

349  
350 One of the GC-MS prenatal diagnostic ratios for PORD remains a challenge for LC-  
351 MS/MS under conditions developed for this study. With fetal PORD there is  
352 increased androsterone production as a result of the “alternative pathway” activity  
353 (Arlt, et al. 2004) resulting in markedly increased androsterone/etiocholanolone  
354 ratio (Shackleton et al 2004a). That ratio should theoretically be determined by  
355 direct analysis of glucuronides and this separation has been already reported by C18  
356 columns both in glucuronides (Pozo et al. 2008) and unconjugated (Marcos and  
357 Pozo, 2016). Unfortunately, under current chromatographic conditions developed  
358 for the disulfates such isobaric monoconjugates (sulfates or glucuronides) could not  
359 be resolved.

#### 360 361 Postnatal detection of PORD:

362  
363 While this paper has focused on prenatal diagnosis of PORD by  $3\beta 5\alpha$ PD-diS  
364 measurement, Shackleton and co-workers (Shackleton et al. 2004b) have shown  
365 that its precursor 5PD-diS is a key analyte in diagnosing the condition in the first  
366 months of life suggested its inclusion here. In PORD neonatal samples this steroid is



dominant, excretory values exceeding the classical major metabolites such as 16 $\alpha$ OHDHEA-S and 16 $\alpha$ -hydroxypregnenolone sulfate whose biosynthesis by 16 $\alpha$ -hydroxylation is also POR dependent. In the first weeks of life the fetal zone of the adrenal is still dominant, but diminishing, and is responsible for producing a large amount of 3 $\beta$ -OH- $\Delta^5$  steroids.

Figure 7 illustrates the separation of steroid disulfates in an affected PORD infant and normal infant. We have included 16 $\alpha$ OHDHEA-S as analyte to act as denominator for a potential diagnostic ratio 5PD-diS/16 $\alpha$ OHDHEA-S. This ratio is shown for three affected infants and normal controls in Figure 7C, clearly defining the condition. Interestingly, one of the first steroid disulfates to be identified in the neonatal period were 5AD(17 $\alpha$  and 17 $\beta$ )-diS (Shackleton, et al. 1968a, Laatikainen, et al. 1972), and 16 $\beta$ OHDHEA-diS (Shackleton, et al. 1968b, Laatikainen, et al. 1972) and these are clearly separated with this methodology (Figure 7).

Smith-Lemli-Opitz Syndrome. "7-dehydrosterol reductase" deficiency. (SLOS) (OMIM 602858 location: 11q13,4)

This condition is caused by deficiency in 7-dehydrosterol reductase and the notable feature is a build-up of 7-and 8-dehydrocholesterol, which can be used to diagnose the condition when measured in amniotic fluid (Kelley,1994). The affected fetus can use these sterols as steroid precursors, resulting in the appearance in maternal urine of dehydro (DH) versions of common natural steroids. For instance, 5 $\beta$ -pregn-7(and 8)-ene-3 $\alpha$ ,17 $\alpha$ ,20S-triol, (7(8)-DHPT) and an estriol equivalent, principally 8-dehydroestriol (8-DHE3) (Guo, et al. 2001, Shackleton, et al. 1999). The biosynthesis of steroids in SLOS pregnancy is illustrated in Figure 8.

SLOS steroids are mainly excreted as glucuronides. Thus, distinct peaks corresponding to different isomers of 8-DHE3-G were found in all SLOS samples (Figure 9). On the other hand, the detection of 7(8)-DHPT-G provided more

difficulties due to endogenous interferences probably coming from other pregnenetriols and DH-hydroxypregnenolones which would share the same transition (Figure 9).

A systematic study has not been made of steroid sulfates in this condition and lack of appropriate authentic compounds has meant that only *candidate* chromatographic peaks were provisionally identified. Such peaks were chosen by having the expected CIL transitions for steroids with additional unsaturation and to be accepted as SLOS-specific “candidate” analytes these peaks had to be present in all six confirmed SLOS pregnancies, and be absent from controls. Two main metabolites were found. Peaks with the expected transitions for DHPT-diS and DHPD-diS were observed in all SLOS samples. Additionally, other minor metabolites such as DH-androstenediol-diS were also found. In Figure 9 the chromatograms of the proposed steroid disulfate analytes with DH-pregnanetriol glucuronide and DHE3-glucuronide are shown. Little information can be stated on stereochemistry of candidate analytes; not only that of 3- and 5-positions but both  $\Delta^7$  and  $\Delta^8$  isomers are likely present. The chromatograms illustrated were from one affected pregnancy and one control. Similar chromatograms were produced for the other five affected pregnancies and controls. Clearly this is the most challenging of the three conditions for conjugate LC-MS/MS analysis although aberrant steroid conjugate peaks definitive for SLOS were detected.

#### *Distinguishing the disorders: summary*

This study has focused on the mass spectrometric analysis of steroid disulfates, but steroid monosulfates and glucuronides have been included where required to determine ratios used for diagnosis. In order to evaluate the potential of the approach based on the combined screening of glucuronides, monosulfates and disulfates, we propose a panel of markers able to differentiate between the selected

disorders and control samples. We found that using the ratios  $16\alpha\text{OHDHEA-S/E3-G}$ ,  $3\beta 5\alpha\text{PD-diS/E3-G}$ ,  $5\text{PD-diS}/16\alpha\text{OHDHEA-S}$ ,  $21\text{OHPreg-diS/E3-G}$ ,  $21\text{OHPreg-diS}/16\alpha\text{OHDHEA-S}$ ,  $8\text{DHE3-G/E3-G}$  and  $\text{DHPT-diS/E3-G}$  allowed for the successful differentiation between the controls and the different disorders. Hopefully the study emphasizes the potential of LC-MS analysis of all conjugate types in future development of steroid metabolomics.

### ***General discussion***

The steroid disulfates are a minor fraction of the urinary steroid metabolome, but may provide significant markers of aberrant steroid biosynthesis. As a family, intact steroid disulfates have not been recently subject to detailed study due a lack of suitable analytical methodology. In the past, studying this family always involved time-consuming fractionation of free and conjugate families followed by solvolysis and GC-MS analysis. Most of the available literature stems from the 60's and 70's and it was shown early which secondary positions (assuming the primary sulfated position is the 3-hydroxyl) could be sulfated. These were 17 ( $\alpha$ - and  $\beta$ ),  $16\beta$ - and 18 in  $\text{C}_{19}$  steroids and 20S- and 21- in  $\text{C}_{21}$  steroids. During that early research period the dominant biological materials chosen to study were associated with pregnancy. In that respect our current studies have followed this lead and the major disulfate components reported here were also noted in the early publications (Shackleton, et al. 1968a; Shackleton, et al. 1968b; Jänne, et al. 1969; Jänne and Vihko 1970; Laatikainen et al 1972; Meng and Sjövall 1997).

There is little definitive evidence as to which sulfotransferases are responsible for the secondary sulfation (Mueller et al 2015), and how disulfates are transported (Grosser et al 2017). Available sulfation evidence points solely to SULT2A1 which appears to have an active site capable of encompassing a wide variety of steroid substrates (both free and monosulfated) and conduct sulfation at either end of the

steroid molecule. Thus, it can sulfate free steroids or steroid monosulfates (Cook, et al. 2009).

A question remains as to whether disulfation is purely a catabolic reaction or if such steroids could be transportable reservoirs of active hormone precursors, as is likely the case for DHEA and estrone sulfates. Guerami and co-workers (1988) have proposed that 21OHPreg-diS is an 11-deoxycorticosterone (DOC) precursor during pregnancy, particularly since circulating levels of this mineralocorticoid and its sulfate are increased during gestation (Corsan, Macdonald and Casey 1997). It is known that the placenta is capable of hydrolyzing 21-sulfates and the enzyme responsible is the usual STS as 21-desulfation does not occur in STSD (Guerami, et al. 1988). Another possible reservoir for disulfates is 5AD(17 $\beta$ )-diS, potentially a testosterone or estradiol precursor. This steroid is also subject to STS action in mammals. In contrast, it is believed that human sulfatases are inactive on 17 $\alpha$ - (C<sub>19</sub> steroids) or 20S--sulfates, a situation shared with the commercial snail and mollusk enzymes used for hydrolysis in steroid analysis (Stevenson, et al. 2014).

In summary, we have provided analytical data on the steroid disulfates through their measurement as intact molecules by LC-MS/MS, employing CIL scan monitoring. We have attempted to use these additional members of the steroid metabolome to distinguish fetal disorders of steroid synthesis. To the best of our knowledge, this is the first time that direct analysis of steroid disulfates has proved its value for clinical diagnosis.

The ultimate goal of these studies is the ability to quantify the whole urinary steroid metabolome as unhydrolyzed conjugates, the monosulfates, disulfates, glucuronides and mixed sulfate-glucuronide conjugates. Studies of the plasma steroid metabolome should also be included. To achieve this goal will require the synthesis of a multitude of authentic steroids including appropriate internal standards and an improvement in chromatographic resolution.

481

**482 Declaration of interest**

483

484 There is no conflict of interest that could be perceived as prejudicing the impartiality  
485 of the research reported

486

**487 Funding**

488

489 Spanish Health National System is acknowledged for OP contract (CPII16/00027).  
490 Strategic Plan for Research and Innovation in Health (PERIS) of the Catalan  
491 government is acknowledge for OK contract (SLT002/16/00007).

492

**493 Acknowledgements**

494 OP wants to acknowledge Alex Gomez-Gomez for his support in the first steps of this  
495 project. CS wants to acknowledge Prof. Wiebke Arlt and her group at IMSR for  
496 continuing collaboration and support in these steroid metabolomic studies. He also  
497 wants to give appreciation to long-term colleagues such as Drs Norman Taylor, Bert  
498 Hauffa, Richard Kelley, Lisa Kratz, and others for early collaborative work on these  
499 topics.

500

501   **REFERENCES**

- 502   Arcos M, Lieberman S 1967 5-Pregnene-3-beta,20-alpha-diol-3-sulfate-20-(2'-  
503   acetamido-2'-deoxy-alpha-D-glucoside) and 5-pregnene-3-beta,20-alpha-diol-  
504   3,20-disulfate. Two novel urinary conjugates. *Biochemistry* 6 2032-2039.
- 505   Arlt W, Walker EA, Draper N, Ivison HE, Ride JP, Hammer F, Chalder SM, Borucka-  
506   Mankiewicz M, Hauffa BP, Malunowicz EM *et al.* 2004 Congenital adrenal  
507   hyperplasia caused by mutant P450 oxidoreductase and human androgen  
508   synthesis: analytical study. *Lancet* 363 2128-2135.
- 509   Cook IT, Duniec-Dmuchowski Z, Kocarek TA, Runge-Morris M, Falany CN 2009 24-  
510   hydroxycholesterol sulfation by human cytosolic sulfotransferases: formation of  
511   monosulfates and disulfates, molecular modeling, sulfatase sensitivity, and  
512   inhibition of liver x receptor activation. *Drug Metabolism & Disposition* 37 2069-  
513   2078.
- 514   Corsan GH, MacDonald PC, Casey ML 1997 Origin of deoxycorticosterone sulfate  
515   (DOC-SO<sub>4</sub>) in plasma of pregnant women: pregnenolone-3,21-disulfate is a  
516   placental precursor of DOC-SO<sub>4</sub>. *The Journal of Steroid Biochemistry and*  
517   *Molecular Biology* 60 331-337.
- 518   Fabregat A, Pozo OJ, Marcos J, Segura J, Ventura R 2013 Use of LC-MS/MS for the  
519   open detection of steroid metabolites conjugated with glucuronic acid. *Analytical*  
520   *Chemistry* 85 5005-5014.
- 521   Galuska CE, Hartmann MF, Sánchez-Guijo A, Bakhaus K, Geyer J, Schuler G,  
522   Zimmer KP, Wudy SA 2013 Profiling intact steroid sulfates and unconjugated  
523   steroids in biological fluids by liquid chromatography-tandem mass spectrometry  
524   (LC-MS-MS). *Analyst* 138 3792-3801.
- 525   Guerami A, Varner MW, Shackleton CH, MacDonald PC, Casey ML 1988 Origin of  
526   deoxycorticosterone and deoxycorticosterone sulfate in human pregnancy:  
527   absence of steroid 21-sulfatase activity in sulfatase-deficient placenta. *Journal of*  
528   *Steroid Biochemistry* 29 57-62.

- 529 Gomes RL, Meredith W, Snape CE, Sephton MA 2009 Analysis of conjugated  
530 steroid androgens: Deconjugation, derivatisation and associated issues. *Journal of*  
531 *Pharmaceutical and Biomedical Analysis* 49 1133-1140.
- 532 Gomez C, Fabregat A, Pozo OJ, Marcos J, Segura J, Ventura R 2014 Analytical  
533 strategies based on mass spectrometric techniques for the study of steroid  
534 metabolism. *TrAC - Trends in Analytical Chemistry* 53 106-116.
- 535 Grosser G, Bennien J, Sánchez-Guijo A, Bakhaus K, Döring B, Hartmann M, Wudy  
536 SA, Geyer J 2017 Transport of steroid 3-sulfates and steroid 17-sulfates by the  
537 sodium-dependent organic anion transporter SOAT (SLC10A6). *The Journal of*  
538 *Steroid Biochemistry and Molecular Biology*. pii: S0960-0760(17)30261-3  
539
- 540 Guo LW, Shackleton CH, Wilson WK 2001 Synthesis of ring B unsaturated estriols.  
541 Confirming the structure of a diagnostic analyte for Smith-Lemli-Opitz syndrome.  
542 *Organic Letters* 3 2547-2550.
- 543 Haddow JE, Palomaki GE, Knight GJ, Cunningham GC, Lustig LS, Boyd PA 1994  
544 Reducing the need for amniocentesis in women 35 years of age or older with  
545 serum markers for screening. *The New England Journal of Medicine* 330 1114-  
546 1118.
- 547 Jänne O, Vihko R, Sjövall J, Sjövall K 1969 Determination of steroid mono- and  
548 disulfates in human plasma. *Clinica Chimica Acta* 23 405-412.
- 549 Jänne O, Vihko R 1970 Neutral steroids in urine during pregnancy. I. Isolation and  
550 identification of dihydroxylated C19 and C21 steroid disulphates. *Acta*  
551 *Endocrinologica* 65 50-68.
- 552 Kelley RI 1995 Diagnosis of Smith-Lemli-Opitz syndrome by gas  
553 chromatography/mass spectrometry of 7-dehydrocholesterol in plasma, amniotic  
554 fluid and cultured skin fibroblasts. *Clinica Chimica Acta* 236 45-58.

- 555 Laatikainen T, Peltonen J, Nylander P 1973 Determination of estriol, estriol  
556 sulfate, progesterone and neutral steroid mono- and disulfates in umbilical cord  
557 blood plasma. *Steroids* 21 347-359.
- 558 Marcos J, Craig WY, Palomaki GE, Kloza EM, Haddow JE, Roberson M, Bradley LA,  
559 Shackleton CH 2009 Maternal urine and serum steroid measurements to identify  
560 steroid sulfatase deficiency (STSD) in second trimester pregnancies. *Prenatal*  
561 *Diagnosis* 29 771-780.
- 562 Marcos J, Renau N, Casals G, Segura J, Ventura R, Pozo OJ 2014 Investigation of  
563 endogenous corticosteroids profiles in human urine based on liquid  
564 chromatography tandem mass spectrometry. *Analytica Chimica Acta* 812 92-104.
- 565 Marcos J, Pozo OJ 2016 Current LC-MS methods and procedures applied to the  
566 identification of new steroid metabolites. *Journal of Steroid Biochemistry and*  
567 *Molecular Biology* 162 41-56
- 568 McLeod MD, Waller CC, Esquivel A, Balcells G, Ventura R, Segura J, Pozo ÓJ 2017  
569 Constant Ion Loss method for the untargeted detection of bis-sulfate metabolites.  
570 *Analytical Chemistry* 89 1602-1609.
- 571 Meng LJ, Sjövall J 1997 Method for combined analysis of profiles of conjugated  
572 progesterone metabolites and bile acids in serum and urine of pregnant women.  
573 *Journal of Chromatography B: Biomedical Sciences and Applications* 688 11-26.
- 574 Mueller JW, Gilligan LC, Idkowiak J, Arlt W, Foster PA 2015 The Regulation of  
575 Steroid Action by Sulfation and Desulfation. *Endocrine Reviews* 36 526-563.
- 576 Pasqualini JR, Jayle MF 1962 Identification of 3beta,21-dihydroxy-5-pregnene-20-  
577 one disulfate in human urine. *The Journal of Clinical Investigation* 41 981-987.
- 578 Pozo OJ, Van Eenoo P, Deventer K, Delbeke FT 2007 Ionization of anabolic steroids  
579 by adduct formation in liquid chromatography electrospray mass spectrometry.  
580 *Journal of Mass Spectrometry* 42 497-516.



- 581 Pozo OJ, Van Eenoo P, Van Thuyne W, Deventer K, Delbeke FT 2008 Direct  
 582 quantification of steroid glucuronides in human urine by liquid chromatography-  
 583 electrospray tandem mass spectrometry. *Journal of Chromatography A* 1183 108-  
 584 118.
- 585 Reisch N, Idkowiak J, Hughes BA, Ivison HE, Abdul-Rahman OA, Hendon LG, Olney  
 586 AH, Nielsen S, Harrison R, Blair EM *et al.* 2013 Prenatal diagnosis of congenital  
 587 adrenal hyperplasia caused by P450 oxidoreductase deficiency. *The Journal of*  
 588 *Clinical Endocrinology & Metabolism* 98 E528-536.
- 589 Shackleton CHL and Marcos P 2006 GC/MS Steroid Profiling: Diagnosis of  
 590 disorders affecting steroid synthesis and metabolism. In *The Encyclopedia of Mass*  
 591 *Spectrometry*, Vol. 8, pp 789-813. Eds M Gross & R Caprioli. Amsterdam: Elsevier.
- 592 Shackleton CH, Kelly RW, Adhikary PM, Brooks CJ, Harkness RA, Sykes PJ, Mitchell  
 593 FL 1968a The identification and measurement of a new steroid 16beta-  
 594 hydroxydehydroepiandrosterone in infant urine. *Steroids* 12 705-716.
- 595 Shackleton CH, Livingstone JR, Mitchell FL 1968b The conjugated 17-hydroxy  
 596 epimers of delta5-androstene-3beta-17-diol in infant and adult urine and  
 597 umbilical cord plasma. *Steroids* 11 299-311.
- 598 Shackleton CH, Straub KM 1982 Direct analysis of steroid conjugates: the use of  
 599 secondary ion mass spectrometry. *Steroids* 40 35-51.
- 600 Shackleton CH 1983 Inborn errors of steroid biosynthesis: detection by a new  
 601 mass-spectrometric method. *Clinical Chemistry* 29 246-249.
- 602 Shackleton CH, Roitman E, Kratz LE, Kelley RI 1999 Midgestational maternal urine  
 603 steroid markers of fetal Smith-Lemli-Opitz (SLO) syndrome (7-dehydrocholesterol  
 604 7-reductase deficiency). *Steroids* 64 446-452.
- 605 Shackleton C, Marcos J, Arlt W, Hauffa BP 2004 Prenatal diagnosis of P450  
 606 oxidoreductase deficiency (ORD): a disorder causing low pregnancy estriol,

607 maternal and fetal virilization, and the Antley-Bixler syndrome phenotype.  
608 American Journal of Medical Genetics Part A 129A 105-112.

609 Shackleton C, Marcos J, Malunowicz EM, Szarras-Czapnik M, Jira P, Taylor NF,  
610 Murphy N, Crushell E, Gottschalk M, Hauffa B *et al.* 2004 Biochemical diagnosis of  
611 Antley-Bixler syndrome by steroid analysis. American Journal of Medical Genetics  
612 Part A 128A 223-231.

613 Shackleton CH, Marcos J, Palomaki GE, Craig WY, Kelley RI, Kratz LE, Haddow JE  
614 2007 Dehydrosteroid measurements in maternal urine or serum for the prenatal  
615 diagnosis of Smith-Lemli-Opitz syndrome (SLOS). American Journal of Medical  
616 Genetics Part A 143A 2129-2136.

617 Stevenson BJ, Waller CC, Ma P, Li K, Cawley AT, Ollis D, McLeod MD 2014  
618 *Pseudomonas aeruginosa* arylsulfatase: a purified enzyme for the mild hydrolysis  
619 of steroid sulfates. Drug Testing and Analysis 7 903-911.

620 Taylor NF, Shackleton CH 1979 Gas chromatographic steroid analysis for diagnosis  
621 of placental sulfatase deficiency: a study of nine patients. The Journal of Clinical  
622 Endocrinology & Metabolism 49 78-86.

623 Tikkanen MJ 1973 Urinary excretion of estriol conjugates in normal pregnancy.  
624 Journal of Steroid Biochemistry 4 57-63.

625 Waller CC, McLeod MD 2014 A simple method for the small scale synthesis and  
626 solid-phase extraction purification of steroid sulfates. Steroids 92 74-80.

627  
628  
629  
630

## FIGURE LEGENDS

### Figure 1

Chromatographic separation obtained for (A) seven synthesized isomers of pregnanediol disulfates and (B) androsterone and etiocholanolone glucuronide. Note that the method optimized for the separation of isomeric disulfate metabolites was not able to separate epimeric glucuronides.

### Figure 2

Steroid synthesis in STSD pregnancy starting from fetal adrenal pregnenolone. Inactivity of STS prevents conversion of  $16\alpha$ OHDHEA-S to estriol in placenta so the former (and its metabolites) is excreted as sulfates by mother.  $C_{21}$  steroid sulfates upstream from  $16\alpha$ OHDHEA-S also pass the placenta without de-sulfation and are directly excreted in maternal urine.

### Figure 3

Selected urinary  $\Delta^5$  steroid sulfate and estriol conjugate analysis in a control and STSD affected pregnancy. Note the markedly increased  $\Delta^5$  steroid mono and disulfates in STSD. Regarding the estriol conjugates it must be noted that glucuronides give lower MS transition responses than sulfates under the reported conditions. In reality, E3-S is a minor metabolite of estriol compared to the 3- and 16-glucuronides, although it appears contrary in the chromatograms.

### Figure 4

Panel A: Ratios of steroid sulfates to E3 glucuronide (3+16) in STSD. The scale represents the ratios of raw peak areas of transitions, not the actual

amount quantified. All ratios show separation of controls from affected pregnancies. Panel B: a measure of the difference between the lowest STSD ratio and highest control ratio. The higher this value, the greater the separation between affected and normal. Two  $\Delta^5$  pregnene di-sulfates are the most efficacious diagnostic analytes. Panel C: combining data from 5PD-diS and 21OHPreg-diS increases discrimination between normal and STSD.

### Figure 5

Steroid biosynthesis and metabolism in PORD and normal pregnancies and neonates. *Normal pregnancy*: fetal adrenal pregnenolone is converted to maternally excreted estriol conjugates (POR essential). *PORD pregnancy*: excess adrenal pregnenolone (due to PORD) is metabolized primarily to maternally excreted  $3\beta 5\alpha$ PD-diS. *Normal neonate*: Excretion product  $16\alpha$ OHDHEA-S and other sulfates. *PORD neonate*: major pregnenolone excretory product 5PD-diS.

### Figure 6

Selected Reaction Monitoring (SRM) transition chromatograms of estriol conjugates and steroid disulfates in control (A) and PORD affected pregnancies (B). Note high excretion of  $3\beta 5\alpha$ PD-diS and 21OHPreg $3\beta 5\alpha$ -diS. (C) Graph shows peak area ratios (analyte/E3-G) for 2 affected pregnancies and 11 controls.

### Figure 7.

SRM chromatograms of control (A) and affected PORD babies (B). The key analyte is the pregnenolone metabolite 5PD-diS and its relative excess is determined by peak area ratio to  $16\alpha$ OHDHEA-S, normally a dominant metabolite in neonates. Discrimination obtained by the use of 5PD-

diS/16 $\alpha$ OHDHEA-S (C) and 21OHPreg-diS/16 $\alpha$ OHDHEA-S (D) between healthy and PORD babies.

**Figure 8.**

Deficiency of 7-dehydrosterol reductase (DHCR7, SLOS) causes 7-or 8-dehydro-cholesterol to be used as fetal precursor for downstream steroids which retain B- ring unsaturation. Dehydro-pregnanetriol (DHPT) and dehydroestriol (DHE3) glucuronides have been used classically for diagnosis but here are candidate disulfates tentatively identified, compounds not seen in controls.

**Figure 9.**

SRM chromatograms of candidate analytes in SLOS pregnancy. (A) control pregnancy and (B) affected pregnancy. Transition chromatograms for known diagnostic steroid glucuronides and candidate disulfates useful for diagnosis. While authentic steroids are not available, these steroids, with appropriate transitions were only present in affected pregnancies. Steroid A/B ring stereochemistry including  $\Delta^7/\Delta^8$  unsaturation is as yet unknown. In the control chromatograms E3 conjugates are shown but all SLOS candidate disulfates and glucuronides are absent.

**Table 1.** SRM parameters of selected steroids

Analyte	Disorder	MW	Retention time (min)	Precursor ion (m/z)	Product ion (m/z)	Collision energy (eV)
<i>Monoconjugates</i>						
E3-3G	all	464	2.3	463	287	30
E3-16G	all	464	9.3	463	287	30
E3-3S	all	368	7.3	367	287	35
16OHDHEA-S	STSD/PORD*	384	16.0	383	97	40
DHE3-G	SLOS	462	8.6/9.1	461	285	30
DHPT-G	SLOS	510	21.3	509	75	30
<i>Disulfates</i>						
5AD(17 $\alpha$ )-diS	STSD	450	15.5	224	351	15
5AD(17 $\beta$ )-diS	STSD	450	13.4	224	351	15
16 $\alpha$ OHDHEA-diS	STSD	464	10.0	231	365	15
16 $\beta$ OHDHEA-diS	STSD	464	8.2	231	365	15
5PT-diS	STSD	494	10.5	246	395	15
21OHPreg-diS	STSD/PORD	492	18.2	245	393	15
5PD-diS	STSD/PORD*	478	16.7	238	379	15
3 $\beta$ 5 $\alpha$ PD-diS	PORD	480	17.6	239	381	15
DH5AD-diS	SLOS	448	11.4	223	349	15
DHPT-diS	SLOS	494	12.6	246	395	15
DHPD-diS	SLOS	478	15.6	238	379	15

\* PORD neonatal




Figure 1




Figure 2






Figure 3




Figure 4




Figure 5




Figure 6




Figure 7




Figure 8




Figure 9.